

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Sorting Category: 05.5 (E)

**Melting of vortex
solid phase in irradiated $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ single crystals in
tilted magnetic fields**

JOVAN MIRKOVIC, Inst of Materials Science, Univ of Tsukuba, 305-8573 Tsukuba, Japan, and Faculty of Sciences, Univ. of Montenegro, Podgorica, Serbia and Montenegro, SERGEY SAVEL'EV, Frontier Research System, The Institute of Physical and Chemical Research (RIKEN), Wako-shi, Saitama, 351-0198, Japan, HIROKAZU SATO, TAKASHI YAMAMOTO, ITSUHIRO KAKEYA, Institute of Materials Science, University of Tsukuba, 305-8573 Tsukuba, Japan, FRANCO NORI, Frontier Research System, The Institute of Physical and Chemical Research (RIKEN), Wako-shi, Saitama, 351-0198, Japan, KAZUO KADOWAKI, Institute of Materials Science, University of Tsukuba, 305-8573 Tsukuba, Japan — The boundary between the vortex-solid and the vortex-liquid in $H_c - H_{ab}$ phase plane for irradiated $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ single crystals is studied by measuring the local ac-magnetic permeability by using the miniature coils in tilted magnetic fields. It was found that the c -axis magnetic field component at the phase transition decreases linearly when increasing the in-plane magnetic field at high temperatures even in a wider angular range than in pristine samples. At lower temperatures, this linear decrease transforms to a concave (hyperbolic-like) curve that differs even stronger from the usual elliptical phase boundary derived from Ginzburg-Landau theory. A theoretical approach to this challenging problem is discussed.

Jovan Mirkovic

jovanmirkovic@cg.yu

Faculty of Sciences, University of Montenegro,

PO Box 211, 81000 Podgorica, Serbia and Montenegro

Prefer Oral Session
 Prefer Poster Session

Date submitted: 04 Dec 2005

Electronic form version 1.4