

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Sorting Category: 17.5 (C)

Study of Atoms and Molecules with Auxiliary-Field Quantum Monte Carlo¹ WIRAWAN PURWANTO, MALLIGA SUEWATTANA², HENRY KRAKAUER, SHIWEI ZHANG, ERIC J. WALTER, College of William and Mary, VA — We study the ground-state properties of second-row atoms and molecules using the phase-less auxiliary-field quantum Monte Carlo (AF QMC) method.³ This method projects the many-body ground state from a trial wave function by means of random walks in the Slater-determinant space. We use a single Slater-determinant trial wave function obtained from density-functional theory (DFT) or Hartree-Fock (HF) calculations. The calculations were done with a plane-wave basis and supercells with periodic boundary condition. We investigate the finite-size effects and the accuracy of pseudopotentials within DFT, HF, and AF QMC frameworks. Pseudopotentials generated from both LDA (OPIUM⁴) and HF⁵ are employed. We find that the many-body QMC calculations show a greater sensitivity to the accuracy of the pseudopotentials. With reliable pseudopotentials, the ionization potentials and dissociation energies obtained using AF QMC are in excellent agreement with the experimental results.

¹Supported by DOE CMSN. Computing was done in CPD and NCSA.

²Currently at Oak Ridge National Lab, TN

³S. Zhang and H. Krakauer, Phys. Rev. Lett. **90**, 136401 (2003)

⁴<http://opium.sourceforge.net>

⁵I. Ovcharenko, A. Aspuru-Guzik, and W. A. Lester, J. Chem. Phys. **114**, 7790 (2001)

Prefer Oral Session
 Prefer Poster Session

Wirawan Purwanto
wxpurw@wm.edu
College of William and Mary

Date submitted: 30 Nov 2005

Electronic form version 1.4