

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Sorting Category: 16.12.3 (E)

Towards 3D and Multilayer Electromagnetic Metamaterials Structures in the THz Range¹ B. D. F. CASSE, Northeastern University, H. O. MOSER, J. W. LEE, S. INGLIS, M. BAHOU, L. K. JIAN, Singapore Synchrotron Light Source — V. G. Veselago predicted that left-handed materials would exhibit a plethora of unusual effects such as a negative index of refraction as used in Snell's law, a reverse Doppler and Čerenkov effect. This novel class of materials, following Pendry's recipes, can potentially restore evanescent waves to focus subwavelength details in an image. Micron-size electromagnetic metamaterials (EM³) structures which exist so far were produced and characterized as single layer structures. Furthermore the structures were produced with a low yield. In the first part of the talk, we will present techniques for producing copious amount of EM³ chips via the LIGA process using Synchrotron radiation and demonstrate assembly of the first multilayer THz EM³ structures. The planar micro- or nanoEM³ produced so far are also highly anisotropic. Recently, we proposed schemes to produce more isotropic structures, within the same matrix, via tilted X-ray exposures that were introduced in the LIGA process years ago. In the second part of the talk, we will show the results of microfabrication of nearly 3D EM³ structures for the THz range.

¹I would like to acknowledge the funding agencies, namely A*STAR and NUS Core Support for financing the project (A*STAR/MOE RP3979908M, A*STAR 0121050038, and NUS Core Support C-380-003-003-001 grants).

- Prefer Oral Session
 Prefer Poster Session

B. D. F. Casse
d.casse@neu.edu
Northeastern University

Date submitted: 20 Nov 2006

Electronic form version 1.4