

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Sorting Category: 13.6.2 (T)

Exact-diagonalization treatment of the non-universal transport regime in few-electron quantum dots¹ LESLIE O. BAKSMATY², CONSTANTINE YANNOULEAS, UZI LANDMAN, School of Physics, Georgia Institute of Technology — Recently, experimental studies³ have revealed a distinct second transport regime in the behavior of transmission phases obtained via Aharonov-Bohm interferometry using small quantum dots (QDs); namely, a non-universal regime for QDs with $N < 10$ electrons, in addition to the earlier known universal one for larger QDs with $N > 14$. Sophisticated (beyond-the-mean-field) many-body methods are needed for describing this non-universal regime. Here, we study the transport properties of small QDs using exact-diagonalization (EXD) calculations in conjunction with Bardeen's theory of quasiparticle mediated conductance.⁴ We will present EXD calculations⁵ for anisotropic QDs, and for a wide range of anisotropies and strengths of inter-electron repulsion.

¹Supported by the U.S. D.O.E. (FG05-86ER-45234)

²Current address: Physics Dept., Rice University

³M. Avinun-Kalish *et al.*, Nature **436**, 529 (2005)

⁴For an adaptation of the formalism to QDs, see S.A. Gurvitz, arXiv:0704.1260v1

⁵C. Yannouleas and U. Landman, Rep. Prog. Phys. **70**, 2067 (2007)

Prefer Oral Session
 Prefer Poster Session

Constantine Yannouleas
constantine.yannouleas@physics.gatech.edu
School of Physics, Georgia Institute of Technology

Date submitted: 26 Nov 2007

Electronic form version 1.4