

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Sorting Category: 13.3 (C)

Few-electron anisotropic quantum dots in low magnetic fields: exact-diagonalization results for excitations, spin configurations, and entanglement¹ CONSTANTINE YANNOULEAS, UZI LANDMAN, School of Physics, Georgia Institute of Technology — Following earlier studies² for $N = 2 - 3$ electrons, exact-diagonalization calculations for $N = 4 - 6$ electrons in anisotropic quantum dots, covering a broad range of confinement anisotropies and strength of inter-electron repulsion, will be presented for zero and low magnetic fields. The excitation spectra are analyzed as a function of the magnetic field and of quantum-dot anisotropy. Analysis of the many-body wave functions through spin-resolved two-point correlations reveals that the electrons tend to localize forming Wigner molecules (WMs). For strong anisotropy, the WMs acquire a linear geometry, and the wave functions with a total spin projection $S_z = (N - 2)/2$ are similar to the strongly entangled W states. For intermediate anisotropy, the WMs exhibit a more complex structure. The degree of entanglement can be quantified through the use of the von Neumann entropy.

¹Supported by the U.S. D.O.E. (FG05-86ER-45234)

²Y. Li, C. Yannouleas, and U. Landman, arXiv:0710.4325v1 [Phys. Rev. B (2007), in press]; C. Yannouleas and U. Landman, Rep. Prog. Phys. **70**, 2067 (2007)

Prefer Oral Session
 Prefer Poster Session

Constantine Yannouleas
constantine.yannouleas@physics.gatech.edu
School of Physics, Georgia Institute of Technology

Date submitted: 26 Nov 2007

Electronic form version 1.4