

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Sorting Category: 15.3 (E)

Dynamical reconstruction of the valence exciton in LiF¹ PETER ABBAMONTE, University of Illinois, WEI KU, Brookhaven National Laboratory, TIM GRABER, University of Chicago, JAMES REED, SERBAN SMADICI, University of Illinois, ABHAY SHUKLA, Universite Pierre et Marie Curie, JEAN-PASCAL RUEFF, Synchrotron SOLIEL — We have used inelastic x-ray scattering, coupled with recently developed inversion techniques, to reconstruct the structure and dynamics of the valence exciton in the prototype alkali halide LiF. Our inversions, which yield resolutions $\Delta x = 0.533\text{\AA}$ and $\Delta t = 20.67\text{as}$ ($2.067 \times 10^{-17}\text{s}$), reveal that the exciton forms in less than 50as , oscillates with a period of 283as , and decays after approximately 5fs . It contains a pronounced $a/3$ internal periodicity, where $a = 4.027\text{\AA}$ is the crystal lattice parameter, that changes little during the course of its life, indicating that this exciton lies very close to the Frenkel limit. Our results resolve a 70 year old controversy about the valence exciton in alkali halides and, when compared to *ab initio* calculations, demonstrate a simplified theoretical approach to describing excitons in the limit of strong binding energy.

¹This work is supported by the US Department of Energy, Office of Basic Energy Sciences, DE-FG02-07ER46459.

Prefer Oral Session
 Prefer Poster Session

Peter Abbamonte
abbamonte@mrl.uiuc.edu
University of Illinois

Date submitted: 01 Dec 2007

Electronic form version 1.4