

Abstract Submitted
for the OSS07 Meeting of
The American Physical Society

Sorting Category: 01 (C)

Resonance sharpening of Aharonov-Bohm (AB) oscillations in an AB ring with double quantum dots¹ ERIC HEDIN, YONG JOE, Ball State University, ARKADY SATANIN, Institute for Physics of Microstructures, Russia — In a balanced AB ring with an identical QD in each arm, a combination of Breit-Wigner and Fano resonances presents a sensitively balanced region of parameter space in which arbitrarily sharp transmission resonance spikes manifest. The transmission becomes a resonance zero when the electron Fermi energy matches the quasi-bound state energy of the matched QD's. However, for values of magnetic flux through the AB ring which are integral multiples of the flux quantum, the analytical expression of the transmission amplitude (obtained from an exactly solvable tight-binding formalism) becomes indeterminate (zero over zero). The resulting transmission resonances near this critical point can show delta-function like full-transmission spikes as a function of magnetic flux. This effect may have practical application as a sensitive monitor of magnetic flux.

¹One of the authors (E. R. H.) is partially supported by a grant from the Center for Energy Research, Education, and Service (CERES) at Ball State University.

Prefer Oral Session
 Prefer Poster Session

Eric Hedin
erhedin@bsu.edu
Ball State University

Date submitted: 16 Apr 2007

Electronic form version 1.4