

Abstract Submitted
for the TSF06 Meeting of
The American Physical Society

Sorting Category: 3.0 (E)

Observation of the high sensitivity of Positron Annihilation induced Auger electron spectroscopy to thermally induced changes in the oxidation state of Cu atoms at the surface of previously oxidized Cu(100)¹ M.P. NADESALINGAM, S. MUKHERJEE, N. FAZLEEY, B.R. DAVIS, J. ZHU, A.H. WEISS, Department of Physics, The University of Texas at Arlington — Changes in the surface of an oxidized Cu(100) single crystal resulting from vacuum annealing over a temperature range from 20 °C to 800 °C has been investigated using Positron annihilation induced Auger electron spectroscopy (PAES). The PAES measurements show a large monotonic increase in the intensity of the annihilation induced Cu ($M_{2,3}VV$) Auger peak as the sample is subjected to a series of isochronal anneals in vacuum up to annealing temperature 300 °C. The intensity then decreases monotonically as the annealing temperature is increase to ~600 °C. The Ps fraction, f_{PS} of these surfaces was found to have the opposite trend going from $f_{PS} = 0.79$ for the surface before any annealing to $f_{PS} = 0.51$ after annealing at 300 °C. These results provide a clear demonstration of the thermal reduction of the copper oxide surface after annealing at 300 °C followed by re-oxidation of the copper surface at the higher annealing temperatures presumably due to the diffusion of subsurface oxygen to the surface.

¹Research is funded by Welch Foundation (grant Y-1100)

Prefer Oral Session
 Prefer Poster Session

Manori Nadesalingam
manori@uta.edu
Department of Physics, The University of Texas at Arlington

Date submitted: 12 Sep 2006

Electronic form version 1.4