

Abstract Submitted
for the NEF14 Meeting of
The American Physical Society

Parameterized tight-binding models of ferromagnetic bilayer manganites M. BAUBLITZ, Boston University, Northeastern University, CHRISTOPHER LANE, Northeastern University, HSIN LIN, Northeastern University, National University of Singapore, HASNAIN HAFIZ, R.S. MARKIEWICZ, B. BARBIELLINI, Northeastern University, Z. SUN, D.S. DESSAU, University of Colorado, A. BANSIL, Northeastern University — Half-metallic behavior in materials has been a subject of extensive research due to its potential for applications in spintronics. Ferromagnetic manganites have been seen as a good candidate, and aside from a small minority-spin pocket observed in $\text{La}_{2-2x}\text{Sr}_{1+2x}\text{Mn}_2\text{O}_7$ ($x = 0.38$), transport measurements show that ferromagnetic manganites essentially behave like half metals. Here we develop robust tight-binding models to describe the electronic band structure of the majority as well as minority spin states of ferromagnetic, spin-canted antiferromagnetic, and fully antiferromagnetic bilayer manganites. Both the bilayer coupling between the MnO_2 planes and the mixing of the $|x^2 - y^2\rangle$ and $|3z^2 - r\rangle$ Mn 3d orbitals play an important role in the subtle behavior of the bilayer splitting. Effects of k_z dispersion are included.

M. Baublitz
Boston University, Northeastern University

Date submitted: 09 Oct 2014

Electronic form version 1.4