

Abstract Submitted
for the OSF08 Meeting of
The American Physical Society

Evidence for Multiple Negative-Parity Band Structure in ^{71}Se

N.R. BAKER, Ohio Wesleyan University, R.A. KAYE, S.R. ARORA, Ohio Wesleyan University, J. BRUCKMAN, Monmouth College, S.L. TABOR, T.A. HINNERS, C.R. HOFFMAN, S. LEE, Florida State University, J. DORING, BFS, Germany — The negative-parity bands of ^{69}Se and ^{73}Se indicate a stark contrast between strong single-particle (^{69}Se) and collective (^{73}Se) behavior over a wide range of spins. However, only one negative-parity band has been observed so far in ^{71}Se , making it difficult to see where it lies between these two very different cases. Thus, the goal of the present work was to extend the level scheme of ^{71}Se as much as possible, with an emphasis on finding new negative-parity states. ^{71}Se nuclei were produced at high spin following the 80-MeV ^{54}Fe (^{23}Na , αpn) reaction at Florida State University. $\gamma - \gamma$ coincidences were measured using an array of 10 Compton-suppressed Ge detectors which included three Clover detectors. From the coincidence relationships, new states were found that formed candidates for perhaps two new negative-parity bands. Cranked-shell model calculations indicate that one new band is associated with rigid-body rotation at high spin.

Robert Kaye
Ohio Wesleyan University

Date submitted: 23 Sep 2008

Electronic form version 1.4