

Abstract Submitted
for the OSF10 Meeting of
The American Physical Society

A Model of Smarandache Geometry in Quantum Mechanics ION PATRASCU, Fratii Buzesti College, Craiova, Romania — Let's consider a simple model of a Smarandache Geometry built in the following way: - an Euclidean plane α , where through any exterior point to a given line (d) there is only one parallel line; - and an Elliptic sphere (S), where lines are defined as the big sphere circles, and points are the regular points on the sphere's surface; this is a Riemannian model of an Elliptic Geometry; - suppose the plane α cuts the sphere (S) upon a big sphere circle (C) into two equal parts; let's A and B be two distinct points on (C), which simultaneously belongs to both: the Euclidean plane α and to the Non-Euclidean sphere (S); therefore, the plane α together with the sphere (S) form a model (M) of a Smarandache Geometry. This model can be interpreted in Quantum Mechanics as follows: - **a particle (P) that it is and it is not in a place in the same time**, is like this circle (C) which is a line [if (C) is referred to the sphere (S)] and it is not a line [if (C) is referred to the plane α] in the model (M) simultaneously; - **a particle (R) which is in two places in the same time**, is like line AB (i.e. the line which passes through the above distinct points A and B) in the model (M); which means that 'line' AB is a straight line in the classical sense in the Euclidean plane α , while 'line' AB is the big sphere circle (C) in the Non-Euclidean sphere (S), therefore line AB is simultaneously in two different places (and has two different forms).

Florentin Smarandache
University of New Mexico, Gallup Campus

Date submitted: 08 Sep 2010

Electronic form version 1.4