

Abstract Submitted
for the OSF20 Meeting of
The American Physical Society

Correlations and dynamic fluctuations in high energy collisions.

ZOULFEKAR MAZLOUM, SEAN GAVIN, Wayne State University, GEORGE MOSCHELLI, Lawrence Technological University, AHMED KHUBRANI, Jazan University — Is thermalization necessary for hydrodynamic flow in nuclear collisions? The discovery of flow-like azimuthal correlations in pA and high-multiplicity pp collisions raises profound questions about the onset of collective flow and its relation to hydrodynamics. We seek independent experimental information on the degree of thermalization in order to identify those hydrodynamic collision systems in which flow is sensitive to equilibrium QCD properties.

We aim to develop a protocol for identifying the degree of thermalization using a combination of momentum and multiplicity correlation. To study the effect of thermalization on these correlations, we use Boltzmann equation in the relaxation time approximation with Langevin noise. We derive a new non-equilibrium transport equation for the two-body distribution function that is consistent with the conservation laws obeyed by microscopic scattering processes. We find that transverse momentum fluctuations in peripheral PbPb collisions at LHC markedly deviate from equilibrium behavior. We propose new measurements that can provide more refined information.

Zoulfekar Mazloun
Wayne State University

Date submitted: 03 Oct 2020

Electronic form version 1.4