

Abstract Submitted
for the OSS05 Meeting of
The American Physical Society

Novel Gapped Quantum Wire TRINANJAN DATTA, ERICA W. CARLSON, Purdue University — High quality state of the art quantum wires (QWRs) can be fabricated by the novel Cleaved Edge Overgrowth (CEO) technique, proposed by (Pfeiffer et al, 1990). Transverse quantization in these QWRs leads to a succession of nested energy bands. With the lowest two successive energy levels occupied, gapped phases are possible including, e.g., an intersubband charge density wave (ICDW) and a Cooper phase with strong superconducting fluctuations (Starykh et al, 2000). Due to the possibility of density reorganization, in which it becomes favorable for the two lowest subbands to match their densities, the ICDW is usually the most likely state. Recently, by exploiting the valley degeneracy in AlAs, a single QWR has been fabricated with two degenerate nonoverlapping bands separated in k space by half an Umklapp vector (Moser et al 2004). For low densities the structure is able to access a multiple subband regime that is not subject to the density reorganizing ICDW, leaving a Cooper phase to flourish. Using Abelian bosonization, we explore the relevant interaction terms for this system, including Umklapp assisted Cooper scattering and discuss the phase diagram.

Trinanjana Datta
Purdue University

Date submitted: 11 Mar 2005

Electronic form version 1.4