

Abstract Submitted
for the OSS12 Meeting of
The American Physical Society

Local structure in hard-sphere chain-molecule fluids¹ SAMBID WASTI, MARK TAYLOR, Dept. of Physics, Hiram College — The conformation of a polymer chain in solvent is coupled to the local structure of the solvent environment. For hard-sphere systems, a monomeric solvent acts to compress a flexible hard-sphere-solute chain and, for a dense system, the local solvent structure is imprinted onto the chain. Here we use Monte Carlo simulation, including bond-rebridging moves, to study the size and conformation of a hard sphere chain in a hard-sphere solvent as a function of both solvent density and solvent diameter. We also study the structure of a hard-sphere-chain solute in a hard-sphere-chain solvent. In the case of a 5-mer chain in 5-mer solvent we show that the effects of solvent can be mapped to a set of two-body solvation potentials. Following our previous work on hard-sphere chains in monomeric solvent [1], we explore the application of these short chain potentials to the study of longer chain-molecule fluids.

[1] M.P. Taylor and S. Ichida, *J. Polym. Sci. B: Polym. Phys.* 45, 3319 (2007).

¹Funding: NSF DMR-0804370

Mark Taylor
Dept. of Physics, Hiram College

Date submitted: 16 Mar 2012

Electronic form version 1.4