

Abstract Submitted  
for the SES05 Meeting of  
The American Physical Society

**Precursors of 1D behavior for  $D > 1$ : evolution of the non-analytic correction to the Fermi-liquid behavior** RONOJOY SAHA, Dept. of Physics, University of Florida, Gainesville, FL-32611-8440, USA, DMITRII MASLOV, Dept. of Physics, University of Florida, Gainesville, Florida 32611-8440, USA — The Fermi-liquid forms of the specific heat ( $C(T)$ ) and static spin susceptibility ( $\chi_s$ ) acquire universal non-analytic corrections[1] and the degree of non-analyticity increase inversely with the dimensionality. This predicts that the strongest non-analyticity in the specific heat should be found in 1D; however, bosonization shows that the  $C(T)$  is analytic in 1D. We resolve this paradox by showing that the general argument, for non-analyticity in  $D > 1$  at the second order in the interaction, breaks down in 1D due to a subtle cancellation and the non-analytic  $T \ln T$  term in the specific heat in 1D occurs at the *third* order for electrons with *spin*. We obtain the same result by considering the RG flow of the marginally irrelevant operator in the sine-Gordon theory. For spinless electrons, the non-analyticities in the particle-particle and particle-hole channels cancel out and the resulting  $C(T)$  is linear in  $T$ . The singularity in the particle-hole channel causes non-analyticity in the spin susceptibility  $\chi_s \propto \ln \max\{|Q|, |H|, T\}$  present at the second order. [1] A.V. Chubukov and D.L. Maslov, Phys. Rev. B 68, 155113 (2003).

Ronojoy Saha

Date submitted: 04 Aug 2005

Electronic form version 1.4