

Abstract Submitted
for the SES07 Meeting of
The American Physical Society

Second harmonic generation in arrays of centrosymmetric gold nano-dimers¹ DAVON W. FERRARA, KEVIN A. TETZ, MATTHEW D. MCMAHON, Vanderbilt University, RICHARD F. HAGLUND, JR. — Second-harmonic generation (SHG) is an important signature of electron dynamics in nanoparticles as well as a sensitive probe of surface effects. In the gap between closely spaced pairs of nanoparticles, or nanodimers (ND), localized electromagnetic field energy creates a “hot spot” that has been shown to affect SHG from asymmetric NDs. We will present new experimental results demonstrating the role that gap size and field localization plays in SHG from centrosymmetric ND arrays. Arrays of ellipsoidal nanoparticles were fabricated using focused ion-beam lithography techniques with gold deposited via thermal evaporation to give particles 20 nm in height with varying areal aspect ratios. In the ND arrays, symmetry forbids SHG in the forward direction, but not at larger angles. Our experiments indicate suppression in SHG intensity with decreasing gap size and evidence of stronger long range interactions between particles with separation over 200 nm. Finite-difference time-domain (FDTD) simulations were also performed in order to correlate field localization with SHG. Our simulations show a strong dependence on the polarization of incident light.

¹Partially supported by Department of Energy grant DE-FG02-01ER45916.

Davon W. Ferrara
Vanderbilt University

Date submitted: 20 Aug 2007

Electronic form version 1.4