

Abstract Submitted  
for the SES08 Meeting of  
The American Physical Society

**The Topology of Chaotic Transport and Escape**<sup>1</sup> JAISON NOVICK,  
College of William and Mary, KEVIN MITCHELL, University of California Merced,  
JOHN DELOS, College of William and Mary — Chaotic transport and escape ap-  
pears in many different systems such as the escape of an asteroid from a planet's  
gravitational field to the escape of ionizing electrons from hydrogen in parallel elec-  
tric and magnetic fields. Numerical simulations have shown that the times to escape  
some region without return possess a complicated fractal structure. These fractals  
result from the intersection of a line of initial conditions and a homoclinic tangle,  
which is formed from the intersections of infinitely long stable and unstable mani-  
folds emanating from an unstable fixed point. Our group has developed Homotopic  
Lobe Dynamics, a topological theory that allows one to predict subsets of the fractals  
seen in numerical simulations. We first show how to apply homotopy to a homo-  
clinic tangle to obtain a set of symbols and a dynamical mapping on the symbols.  
A symbol and its mappings encode the evolution of an entire family of trajectories.  
Given a symbol and its mappings, we show how to construct a theoretical fractal.  
Finally, we compare a predicted fractal to one obtained from a numerical simulation  
of trajectories propagating in an open chaotic vase-shaped billiard.

<sup>1</sup>Supported by NSF

Jaison Novick  
College of William and Mary

Date submitted: 14 Aug 2008

Electronic form version 1.4