

SES11-2011-000067

Abstract for an Invited Paper
for the SES11 Meeting of
the American Physical Society

The Emergence of Community Structure in Metacommunities¹

PER ARNE RIKVOLD, Department of Physics, Florida State University

The role of space in determining species coexistence and community structure is well established. However, previous studies mainly focus on simple competition and predation systems, and the role of mutualistic interspecies interactions is not well understood. Here we use a spatially explicit metacommunity model, in which new species enter by a mutation process, to study the effect of fitness-dependent dispersal on the structure of communities with interactions comprising mutualism, competition, and exploitation [1,2]. We find that the diversity and the structure of the interaction network undergo a nonequilibrium phase transition with increasing dispersal rate. *Low* dispersion rate favors spontaneous emergence of many dissimilar, strongly mutualistic and species-poor local communities. Due to the local dissimilarities, the global diversity is high. *High* dispersion rate promotes local biodiversity and supports similar, species-rich local communities with a wide range of interactions. The strong similarity between neighboring local communities leads to reduced global diversity.

[1] E. Filotas, M. Grant, L. Parrott, P.A. Rikvold, J. Theor. Biol. **266**, 419 (2010).

[2] E. Filotas, M. Grant, L. Parrott, P.A. Rikvold, Ecol. Modell. **221**, 885 (2010).

¹Supported by U.S. NSF, SERC of Canada, and le Fond Québécois de la Recherche sur la Nature et les Technologies.