

Abstract Submitted
for the SES16 Meeting of
The American Physical Society

Graphene quantum dot synthesis using nanosecond laser pulses¹ KHOMIDKHODZHA KHOLIKOV, DOVLETGELDI SEYTTLIYEV, DANA BIECHELE-SPEZIALE, ZACHARY THOMAS, SKYLER SMITH, SELAHADDIN GUMUS, BENJAMIN KASH, ALI ER, Western Kentucky Univ — A biocompatible photodynamic therapy agent that generates a high amount of singlet oxygen with high water dispersibility and excellent photostability is desirable. In this work, a graphene based biomaterial which is a promising alternative to a standard photosensitizers was produced. Methylene blue was used as a reference photosensitizer. Bacteria deactivation by methylene blue was shown to be inhibited inside human blood due to protein binding. Graphene quantum dots (GQD) were synthesized by irradiating benzene and nickel oxide mixture using nanosecond laser pulses. High resolution transmission electron microscopy (HR-TEM), scanning electron microscopy (SEM), atomic force microscopy (AFM), Fourier transform infrared (FTIR) spectroscopy, and nuclear magnetic resonance (NMR) were used for characterization of GQDs. Initial results show graphene quantum dots whose size less than 10 nm were successfully obtained. UV-VIS spectra shows absorption peak around 310 nm. The results of these studies can potentially be used to develop therapies for the eradication of pathogens in open wounds, burns, or skin cancers. New therapies for these conditions are particularly needed when antibiotic-resistant infections are present.

¹Kentucky Biomedical Research Infrastructure and INBRE

Dovletgeldi Seyitliyev
Western Kentucky Univ

Date submitted: 07 Oct 2016

Electronic form version 1.4