

Abstract Submitted
for the SHOCK11 Meeting of
The American Physical Society

Dynamic Behaviors of Lead Flyer Driven by Collision of Head-on Sliding Detonations CHONGYU ZHANG, Institute of Fluid Physics, CAEP, Mianyang, HAIBO HU, QINGZHONG LI, ZHENGTAO ZHANG, XUELIN SUN, LABORATORY FOR SHOCK WAVES AND DETONATION PHYSICS RESEARCH TEAM — The dynamic behaviors of lead plate driven by head-on sliding detonation waves were characterized with the help of high-speed frame photography and pulsed X-ray radiography. Experimental records have shown a jet like bulging in the collision region, size of which extended rapidly after the collision of the head-on detonation waves because of the obvious speed gradients of particles inside the bulging from the tip to the bottom of the bulging. Multi-layer like structure of loading front formed in the result of the impact of two symmetric detonation fronts. The mass densities inside the bulging structure fixed by the pulsed X-ray radiography were evaluated at the level of 1%~10% from the initial density of lead. The dynamic strength and shock wave melting should have played dominate role in the formation of the initial stage and the evolution of cavitations and fragmentation process finished merely in microseconds inside the continuum of melted lead under the intensive tension of release wave, in the result of which a porous or dispersed stage bulging was formed.

Chongyu Zhang
Institute of Fluid Physics, CAEP, Mianyang

Date submitted: 31 Jan 2011

Electronic form version 1.4