

Abstract Submitted
for the 4CF09 Meeting of
The American Physical Society

Polarizability of Pb III from spectroscopy of high-L Rydberg states of Pb II¹ MARK E. HANNI, JULIE A. KEELE, STEPHEN R. LUNDEEN, Colorado State University, WILLIAM G. STURRUS, Youngstown State University, CHARLES W. FEHRENBACH, Kansas State University, RESIS COLLABORATION — Using the Resonant Excitation Stark Ionization Spectroscopy technique[1], we measured resolved fine structure components of the $n=20$ to $n'=52$ transition in Pb^+ , and determined the polarizability of Pb^{2+} . A critical part of the measurement consists of the determination of the L-value of one or more of the resolved excitation peaks. These measurements were motivated by a discrepancy between previous polarizability determinations based on contrasting methods[2].

[1] S.R. Lundeen and C.W. Fehrenbach, Phys. Rev. A 75, 032523 (2007)

[2] Nicholas Reshetnikov, et. al., Physica Scripta 77, 015301 (2008).

¹Supported by the Chemical Sciences, Geosciences, and Biosciences Division of the Office of Basic Energy Science, U.S. Department of Energy.

Mark E. Hanni
Colorado State University

Date submitted: 28 Sep 2009

Electronic form version 1.4