

Abstract Submitted
for the APR07 Meeting of
The American Physical Society

Advances in Black-Hole Mergers: Spins and Unequal Masses¹

BERNARD KELLY, JAMES VAN METER, NASA Goddard Space Flight Center, SEAN MCWILLIAMS, University of Maryland, DAE-IL CHOI, Korea Institute of Science and Technology Information, JOAN CENTRELLA, NASA Goddard Space Flight Center, WILLIAM DARIAN BOGGS, University of Maryland, JOHN BAKER, NASA Goddard Space Flight Center — The last two years have seen incredible development in numerical relativity: from fractions of an orbit, evolutions of an equal-mass binary have reached multiple orbits, and convergent gravitational waveforms have been produced from several research groups and numerical codes. We are now able to move our attention from pure numerics to astrophysics, and address scenarios relevant to current and future gravitational-wave detectors. Over the last 12 months at NASA Goddard, we have extended the accuracy of our Hahn-Dol code, and used it to move toward these goals. We have achieved high-accuracy simulations of black-hole binaries of low initial eccentricity, with enough orbits of inspiral before merger to allow us to produce hybrid waveforms that reflect accurately the entire lifetime of the BH binary. We are extending this work, looking at the effects of unequal masses and spins.

¹Supported in part by NASA grant O5-BEFS-05-0044

Bernard Kelly
NASA Goddard Space Flight Center

Date submitted: 12 Jan 2007

Electronic form version 1.4