

Abstract Submitted
for the APR09 Meeting of
The American Physical Society

Ion heating due to reconnection and turbulence in SSX¹ T.G. GRAY, M.R. BROWN, Swarthmore College — Ion heating is measured in the SSX device with a high resolution ion Doppler spectrometer. The SSX IDS instrument measures the width and Doppler shift of either the nascent C_{III} impurity 229.7 nm line or a doped He_{II} impurity 468.6 nm line to determine the temperature and line-averaged flow velocity. The velocity resolution of the instrument is ≤ 5 km/s. There is enough signal to resolve the full line within an MHD dynamical time (about 1 μ s in SSX). Spheromak merging in a new oblate flux conserver ($R = 0.25$ m, $L = 0.4$ m) has resulted in some stable configurations but often results in excitation of several unstable MHD modes. After reconnection and instability, we measure a period of ion heating with peak temperatures for carbon $T_C = 40$ eV but for helium only $T_{He} = 20$ eV. Our results are consistent with temperature increasing with ion mass but only scaling like $\sqrt{M_i/m_p}$. We plan a more comprehensive study of the effect of reconnection and turbulence on ion heating for various ion masses (He, C, Xe). A new high purity gas delivery and mixing system is under construction. Results of ion heating with other ions will be presented.

¹Supported by US DOE and NSF.

T. G. Gray
Swarthmore College

Date submitted: 25 Mar 2009

Electronic form version 1.4