

Abstract Submitted  
for the APR10 Meeting of  
The American Physical Society

**Plans to Search for New Particles Decaying to Dijets at CMS**

SERTAC OZTURK, University of Cukurova / Fermilab, ROBERT HARRIS, Fermilab — The Compact Muon Solenoid (CMS) is one of two multi-purpose detectors which is located on the Large Hadron Collider (LHC) at CERN. We use a generated pseudo-data sample corresponding to  $10 \text{ pb}^{-1}$  of integrated luminosity at a pp collision energy of 10 TeV to test CMS plans to search for new particles decaying to two high energy jets (dijets). By construction the measured dijet mass spectrum agrees with the QCD prediction. Applying our search to this pseudo-data sample, we set upper limits on the cross section for dijet resonances. This analysis shows that CMS will be sensitive to new particles decaying to dijets, and if we measured this pseudo-dataset produced from QCD alone, we would exclude at 95% confidence level the following new particles: axiglons and flavor universal colorons with mass below  $1.8 \text{ TeV}$ , excited quarks with mass below  $1.8 \text{ TeV}$  and  $E_6$  diquarks with mass below  $1.0 \text{ TeV}$  and mass between  $1.3 \text{ TeV}$  and  $1.7 \text{ TeV}$ .

Sertac Ozturk  
University of Cukurova / Fermilab

Date submitted: 15 Oct 2009

Electronic form version 1.4