

Abstract Submitted
for the APR16 Meeting of
The American Physical Society

Quasi-collisional Magneto-optic Effects in Collisionless Plasmas

BRETT KEENAN, ALEX FORD, MIKHAIL MEDVEDEV, Univ of Kansas — High-amplitude, chaotic/turbulent electromagnetic fluctuations are ubiquitous in astrophysical plasmas, where they can be excited by various kinetic-streaming and/or anisotropy-driven instabilities, such as the Weibel instability. These fields typically exist on “sub-Larmor scales” — scales smaller than the electron Larmor radius. Electrons moving through such magnetic fields undergo small-angle stochastic deflections of their pitch-angles, thus establishing diffusive transport on long time-scales. We show that this behavior, under certain conditions, is equivalent to Coulomb collisions in collisional plasmas. The magnetic pitch-angle diffusion coefficient, which acts as an effective “collision” frequency, may be substantial in these, otherwise, collisionless environments. We show that this effect, colloquially referred to as the plasma “quasi-collisionality”, may radically alter the expected radiative transport properties of candidate plasmas. We argue that the modified magneto-optic effects in these plasmas provide an attractive, novel radiative diagnostic tool for the exploration and characterization of small-scale magnetic turbulence.

Mikhail Medvedev
Univ of Kansas

Date submitted: 08 Jan 2016

Electronic form version 1.4