

Abstract Submitted
for the APR18 Meeting of
The American Physical Society

Method for robust extraction of proton charge radius from electron-proton-scattering experiments¹ XUEFEI YAN, Duke Univ, PRAD COLLABORATION — A lot of efforts have been devoted to the measurements of the radius of the proton charge distribution (R), but results from different experiments and/or analyses have sizable differences. In high-precision muonic hydrogen Lamb shift experiments, R was measured to be 0.8409 ± 0.0004 fm, while typical global data analyses of the electron-proton (ep) scattering experiments yielded $R = 0.879 \pm 0.009$ fm. This difference has been known as the proton radius puzzle. Some recent global ep -scattering-data analyses found $R \approx 0.84$ fm, but the fitting methods used by them were criticized by others in the literature. We formulated a robust fitting method to find the “real” R in upcoming ep -scattering experiments. Comprehensive tests have been carried out using the binning and statistical uncertainties of the PRad experiment (E12-11-106²). In this talk, we will discuss the details of this method and present the test results.

¹This work is supported in part by the U.S. Department of Energy under Contract No. DE-FG02-03ER41231, NSF MRI award PHY-1229153, Thomas Jefferson National Accelerator Facility and Duke University

²Spokespersons: A. Gasparian (contact), H. Gao, M. Khandaker, D. Dutta

XUEFEI YAN
Duke Univ

Date submitted: 12 Jan 2018

Electronic form version 1.4