

Abstract Submitted  
for the APR20 Meeting of  
The American Physical Society

**The BeEST Experiment: A Search for keV-Scale Neutrinos in the EC Decay of  ${}^7\text{Be}$  with Superconducting Quantum Sensors** KYLE LEACH, Colorado School of Mines, STEPHAN FRIEDRICH, Lawrence Livermore National Laboratory, BEEST COLLABORATION — The search for sterile neutrinos is among the brightest possibilities in our quest for understanding the microscopic nature of dark matter in our universe. Sterile neutrinos - unlike the active neutrinos in the SM - do not interact with normal matter as they move through space, and as such must be observed via their mass-generated effects that result from momentum conservation with SM particles. One way to observe these momentum recoil effects experimentally is through high-precision measurements of electron-capture (EC) nuclear decay, where the final state only contains the neutrino and a recoiling atom. This approach is the most powerful method for BSM neutrino mass searches since it relies only on the existence of a heavy neutrino admixture to the active neutrinos, which is a generic feature of neutrino mass mechanisms, and not on the model-dependent details of their interactions. In this talk, we report the first measurements in the Beryllium EC STJ (BeEST) experimental program, which uses the decay-momentum reconstruction technique to precisely measure the  ${}^7\text{Be}\rightarrow{}^7\text{Li}$  recoil spectrum via  ${}^7\text{Be}$  ions implanted into sensitive superconducting tunnel junction (STJ) radiation detectors.

Kyle Leach  
Colorado School of Mines

Date submitted: 14 Jan 2020

Electronic form version 1.4