

Abstract Submitted
for the APR21 Meeting of
The American Physical Society

Limits on Intrinsic Charm Production from the SeaQuest Experiment¹ RAMONA VOGT, Lawrence Livermore Natl Lab and UC Davis — A nonperturbative charm production contribution, known as intrinsic charm, has long been speculated but has never been satisfactorily proven. The SeaQuest experiment at FNAL is in an ideal kinematic region to provide evidence of J/ψ production by intrinsic charm. Here, J/ψ production in the SeaQuest kinematics is calculated with a combination of perturbative QCD and intrinsic charm to see whether the SeaQuest data can put limits on an intrinsic charm contribution². J/ψ production in perturbative QCD is calculated to next-to-leading order in the cross section. Cold nuclear matter effects included in this component are nuclear modification of the parton densities, absorption by nucleons, and p_T broadening by multiple scattering. The J/ψ contribution from intrinsic charm is calculated assuming production from a $|uudc\bar{c}\rangle$ Fock state. The nuclear modification factor, R_{pA} , is calculated as a function of x_F and p_T for $p + C$, $p + Fe$, and $p + W$ interactions relative to $p + d$. It is shown that the SeaQuest kinematic acceptance is ideal for setting limits on intrinsic charm in the proton.

¹This work was performed under the auspices of the U.S. DoE by LLNL under Contract DE-AC52-07NA27344 and supported by LDRD project 21-LW-034.

²R. Vogt, to be submitted to Phys. Rev. C.

Ramona Vogt
Lawrence Livermore Natl Lab and UC Davis

Date submitted: 16 Dec 2020

Electronic form version 1.4