

Abstract Submitted
for the APR21 Meeting of
The American Physical Society

Measurement of proton-Xi correlation function in Au+Au Collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV by STAR experiment at RHIC KE MI, Central China Normal University — The study of baryon-baryon interactions are important to understand existence of strangelets in heavy-ion collisions and various exotic hadrons, modeling of astronomical objects such as neutron stars. A detailed knowledge of nucleon-nucleon (NN) potentials exists in literature, however very little is known about interactions between anti-nucleons. Similarly lack of scattering data for hyperon-nucleon (YN) systems makes it difficult to construct YN potentials. In heavy-ion collisions, a large number of baryons are produced in each nucleus-nucleus collision, which allows us to study the NN and YN interactions. Measurements of two-particle correlation function are used to study the space-time dynamics of the source created in heavy-ion collisions. At low relative momentum, the two-particle correlations are effected by the Final State Interactions (FSI), making it possible to measure FSI between nucleon and multi-strange baryon Xi. The first measurement of proton-Xi correlation function from the STAR experiment at RHIC for the Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV will be presented.

Ke Mi
Central China Normal University

Date submitted: 09 Jan 2021

Electronic form version 1.4