

Abstract Submitted
for the DAMOP07 Meeting of
The American Physical Society

Complete Single-Center Basis Sets in Atomic Calculations¹

SCOTT I. YOUNG, KYLE ROLLIN, MICHAEL W.J. BROMLEY, Department of Physics, San Diego State University, JIM MITROY, Faculty of Technology, Charles Darwin University, KURUNATHAN RATNAVELU, Faculty of Science, Universiti Malaya, Kuala Lumpur — Single particle orbitals centered on the nucleus are the most commonly used basis in large-scale calculations of atomic structure. The convergence towards a complete basis set, with respect to both the number and angular momenta of the orbitals included in a configuration interaction (CI) expansion, has been investigated using the ground and excited states of the helium atom [1,2,3]. This enabled energies to be determined to within 10^{-8} Hartree, whilst the convergence of the electron-electron δ -function and other relativistic corrections have been examined in detail. Unusual convergence patterns in the CI-Kohn variational scattering method have been observed and, however, high-precision calculations of elastic positron scattering from atoms are achievable [4].

[1] J.Mitroy, M.W.J.Bromley, K.Ratnavelu *Int.J.Quant.Chem.* **107** 907 (2007).

[2] M.W.J.Bromley and J.Mitroy *Int.J.Quant.Chem.* **107** 1150 (2007).

[3] K.Rollin, M.W.J.Bromley and J.Mitroy (in preparation).

[4] S.I.Young, M.W.J.Bromley and J.Mitroy (in preparation).

¹Supported in part by the Australian Research Council.

Michael W.J. Bromley
Department of Physics, San Diego State University

Date submitted: 02 Feb 2007

Electronic form version 1.4