

Abstract Submitted  
for the DAMOP07 Meeting of  
The American Physical Society

**Optimal phase estimation for a Mach-Zehnder interferometer.**

LUCA PEZZE', AUGUSTO SMERZI, BEC-CNR-INFN and Dipartimento di Fisica, Universita' di Trento, I-38050 Povo, Italy, GEORGE KHOURY, JUAN HODELIN, DIRK BOUWMEESTER, Department of Physics, University of California, Santa Barbara, California 93106, USA — We study a Mach-Zehnder interferometer fed by a coherent state in one input port and vacuum in the other. We explore a Bayesian phase estimation strategy to demonstrate that it is possible to achieve the standard quantum limit \*independently\* from the true value of the phase shift and specific assumptions on the noise of the interferometer. We have been able to implement the protocol using parallel operation of two photon-number-resolving detectors and multiphoton coincidence logic electronics at the output ports of a weakly-illuminated Mach-Zehnder interferometer. This protocol is unbiased and saturates the Cramer-Rao phase uncertainty bound and, therefore, is an optimal phase estimation strategy.

Luca Pezze'  
BEC-CNR-INFN and Dipartimento di Fisica, Universita' di Trento, I-38050 Povo, Italy

Date submitted: 08 Mar 2007

Electronic form version 1.4