

Abstract Submitted  
for the DAMOP08 Meeting of  
The American Physical Society

**Magnetic Trapping of Atomic Nitrogen and Cotrapping of NH**

MATTHEW HUMMON, WESLEY CAMPBELL, Department of Physics, Harvard University; Harvard-MIT Center for Ultracold Atoms, HSIN-I LU, Division of Engineering and Applied Sciences, Harvard University; Harvard-MIT Center for Ultracold Atoms, EDEM TSIKATA, YIHUA WANG, JOHN DOYLE, Department of Physics, Harvard University; Harvard-MIT Center for Ultracold Atoms — We observe magnetic trapping of atomic nitrogen ( $^{14}\text{N}$ ) and cotrapping of  $^{14}\text{NH}$  ( $X^3\Sigma^-$ ). We use buffer gas cooling to load the magnetic trap directly from a room temperature molecular beam generated by a radio-frequency plasma source. We trap approximately  $1 \times 10^{11}$   $^{14}\text{N}$  atoms at a peak density of  $5 \times 10^{11} \text{ cm}^{-3}$  at a temperature of  $\approx 550$  mK. The  $1/e$  lifetime of nitrogen in the trap of  $12 +5/-3$  s is limited by collisions with the helium buffer gas. This lifetime sets a limit on the inelastic rate constant for  $^{14}\text{N}-^3\text{He}$  collisions of  $\Gamma_{in} < 2.2 \times 10^{-16} \text{ cm}^3 \text{ s}^{-1}$ .  $^{14}\text{N}$  and  $^{14}\text{NH}$  are cotrapped, with  $4 \times 10^{10}$   $^{14}\text{N}$  atoms and  $1 \times 10^8$   $^{14}\text{NH}$  molecules at peak densities of  $n_N \approx 1 \times 10^{11} \text{ cm}^{-3}$  and  $n_{NH} \approx 1 \times 10^8$  at a temperature of  $\approx 550$  mK.

Matthew Hummon  
Department of Physics, Harvard University

Date submitted: 31 Jan 2008

Electronic form version 1.4