

Abstract Submitted
for the DAMOP08 Meeting of
The American Physical Society

Atomic rubidium, the workhorse of theoretical collision physics

BOUDEWIJN VERHAAR, Eindhoven University, ERIC VAN KEMPEN, Philips Applied Technology, Eindhoven, SERVAAS KOKKELMANS, Eindhoven University, COHERENCE AND QUANTUM TECHNOLOGY TEAM — Since the first realizations of Bose-Einstein condensates in ultracold atomic gases in 1995, the 85Rb and 87Rb atomic species have acted as the workhorses of experimental developments in this field. Parallel to and partly preceding this work the same isotopes figured also as workhorses for successful theoretical attempts aiming at unravelling the needed information on interaction properties like scattering lengths and Feshbach resonances¹. The purpose of this contribution is to describe recent developments of this theoretical work, based on an adiabatic variant of the accumulated phase method, which gives rise to the most reliable and accurate predictions of interaction properties available to date for any alkali atomic gas². The method can be readily applied to predict interaction properties between two different atomic species, once a limited set of experimental data is available to determine the accumulated phases.

¹Hugo Boesten, PhD thesis Eindhoven University (1996, ISBN 90-386-0019-4)

²See preliminary description by E. van Kempen, S. Kokkelmans, D. Heinzen, and B. Verhaar, Phys. Rev. Lett. 88, 093201 (2002).

Boudewijn Verhaar
Eindhoven University of Technology

Date submitted: 01 Feb 2008

Electronic form version 1.4