

Abstract Submitted
for the DAMOP12 Meeting of
The American Physical Society

Strongly interacting fermions in a 1D optical lattice¹ ARIEL SOMMER, LAWRENCE CHEUK, MARK KU, WASEEM BAKR, TARIK YEFSAH, MARTIN ZWIERLEIN, Department of Physics, MIT-Harvard Center for Ultracold Atoms, and Research Laboratory of Electronics, MIT, Cambridge, Massachusetts 02139, USA — Strongly correlated fermions in an array of two-dimensional planes coupled via tunneling serve as an important model system for high-temperature superconductors and layered organic conductors. We realize this model using ultracold fermionic ⁶Li atoms in a one-dimensional optical lattice near a Feshbach resonance. The depth of the lattice controls the interlayer coupling, and tunes the system between three and two dimensions. Pairing between fermions is studied using radio-frequency spectroscopy. The binding energy of fermion pairs is determined along the dimensional crossover and for different interaction strengths through the BEC-BCS crossover.

¹This work was supported by the NSF, AFOSR-MURI, ARO-MURI, ONR, DARPA YFA, a grant from the Army Research Office with funding from the DARPA OLE program, the David and Lucile Packard Foundation and the Alfred P. Sloan Foundation

Ariel Sommer
Department of Physics, MIT-Harvard Center for Ultracold Atoms,
and Research Laboratory of Electronics, MIT,
Cambridge, Massachusetts 02139, USA

Date submitted: 30 Jan 2012

Electronic form version 1.4