

Abstract Submitted
for the DAMOP15 Meeting of
The American Physical Society

Laser cooling and trapping with optical frequency combs

XUEPING LONG, ANDREW JAYICH, ANTHONY RANSFORD, ANNA WANG, WESLEY CAMPBELL, Univ of California - Los Angeles — A large number of atoms and molecules are difficult to control with continuous wave lasers because generating sufficient power at all of the necessary wavelengths is technologically challenging. Mode-locked lasers, through their enhanced efficiency of nonlinear frequency conversion, provide some of these hard to access wavelengths. As a step towards control of exotic atoms and molecules we report on laser cooling and trapping of atoms using an optical frequency comb in two different regimes. Using a single comb, we have created a simultaneous dual-species (isotopes) MOT, demonstrating that multiple comb teeth can be used in parallel to cool and confine species requiring many cw lasers. Separately, we demonstrate comb-based laser cooling on a two-photon transition, which efficiently uses the full time-averaged optical power of the entire comb [1]. Our progress toward extending this to include trapping by making a MOT using this two-photon transition is presented. This work is supported by the National Science Foundation

[1] D. Kielpinski, Phys. Rev. A 73, 063407 (2006)

Long Xueping
Univ of California - Los Angeles

Date submitted: 30 Jan 2015

Electronic form version 1.4