

Abstract Submitted  
for the DAMOP16 Meeting of  
The American Physical Society

**Nuclear Spin Dependent Parity Violation in Diatomic Molecules**

EMINE ALTUNTAS, SIDNEY CAHN, DAVID DEMILLE, Yale University, MIKHAIL KOZLOV, Petersburg Nuclear Physics Institute — Nuclear spin-dependent parity violation (NSD-PV) effects arise from exchange of the  $Z^0$  boson between electrons and the nucleus, and from interaction of electrons with the nuclear anapole moment, a parity-odd magnetic moment. The latter scales with nucleon number of the nucleus  $A$  as  $A^{2/3}$ , whereas the  $Z^0$  coupling is independent of  $A$ . Thus the former is the dominant source of NSD-PV for nuclei with  $A \geq 20$ . We study NSD-PV effects using diatomic molecules, where signals are dramatically amplified by bringing rotational levels of opposite parity close to degeneracy in a strong magnetic field. The NSD-PV interaction matrix element is measured using a Stark-interference technique. We present results that demonstrate statistical sensitivity to NSD-PV effects surpassing that of any previous atomic parity violation measurement, using the test system  $^{138}\text{Ba}^{19}\text{F}$ . We report our progress on measuring and cancelling systematic effects due to combination of non-reversing stray  $E$ -fields,  $E_{nr}$  with  $B$ -field inhomogeneities. Short-term prospects for measuring the nuclear anapole moment of  $^{137}\text{Ba}^{19}\text{F}$  are discussed. In the long term, our technique is sufficiently general and sensitive to enable measurements across a broad range of nuclei.

emine altuntas  
yale university

Date submitted: 28 Jan 2016

Electronic form version 1.4