

Abstract Submitted  
for the DAMOP20 Meeting of  
The American Physical Society

**Progress towards Ultrasensitive Gravity Gradiometers using Macroscopically Delocalized Strontium**<sup>1</sup> KENNETH DEROSE, JAYAMPATHI KANGARA, NATASHA SACHDEVA, TEJAS DESHPANDE, YIPING WANG, JONAH GLICK, TIMOTHY KOVACHY, Northwestern University — Recent precision measurements based on torsion balances and pendulums disagree on the gravitational constant  $G$  by nearly 40 times the smallest reported uncertainty [1,2]. To address this discrepancy, it is important to measure  $G$  with a variety of different methods. Quantum sensors based on atom interferometry have proven to be a powerful tool for measuring  $G$ , with a different set of systematic errors than the classical techniques used in most measurements [2]. Here, we discuss progress toward a new atom interferometric measurement of  $G$  that will leverage recent advances in ultrasensitive atomic gravity gradiometers. We will detail our designs and progress toward the construction of a 2 m fountain capable of delocalizing atomic wavefunctions on a macroscopic scale by utilizing recent advances in large momentum transfer on the strontium transitions. We intend to test our gravity gradiometer with two large single-crystal silicon proof masses. The masses will translate on thick, level granite slabs between measurements where a high-resolution atomic phase readout will allow the determination of  $G$ . In addition, the apparatus will be used to test the gravitational inverse square law in order to search for new particles beyond the standard model. [1] A. Mann. PNAS 113, 9949-9952 (2016); Q. Li et al., Nature 560, 582-588 (2018). [2] G. Rosi, F. Sorrentino, L. Cacciapuoti, M. Prevedeli, and G. M. Tino, Nature 510, 518 (2014).

<sup>1</sup>Office of Naval Research, NIST

Kenneth DeRose  
Northwestern University

Date submitted: 31 Jan 2020

Electronic form version 1.4