

Abstract Submitted
for the DFD10 Meeting of
The American Physical Society

Temporal behavior and vortex topology of topographic wave-breaking OLIVIER EIFF, U. Toulouse, CNRS-IMFT, NICOLAS BOULANGER, KARINE LEROUX, ALEXANDRE PACI, Meteo-France/CNRM, CNRS-GAME — At low Froude numbers or strong stratification, the internal waves generated by flow over an obstacle or mountain will overturn and break. Surprisingly little is known, however, of the dynamics of the wave breaking itself. Afanasyev and Peltier (1998) investigated the wave breaking region via LES and Eiff et al. (2005) via PIV measurements, but both presumed a statistically stationary wave-breaking process after the initial wave overturning. Here, we propose to take a closer look at this assumption by closely analyzing the spatio-temporal structure of internal wave breaking region and the surrounding flow. The analysis is based on Hovmöller diagrams and spatial correlations obtained from 2D-PIV measurements of flows generated in uniform stratified flow over 2D and quasi-2D obstacles in salt-stratified hydraulic channels at different Reynolds numbers ranging from laminar to turbulent. The results reveal low frequency variations throughout the flow field, in and outside the wave-breaking region. This characteristic frequency can be related to be due to a sequence of growth and decay of wave-breaking. Finally, new 3D-3C PIV measurements at high Reynolds numbers reveal a first glimpse of the 3D vortex topology.

Olivier Eiff
U. Toulouse, CNRS-IMFT

Date submitted: 02 Aug 2010

Electronic form version 1.4