

Abstract Submitted
for the DFD10 Meeting of
The American Physical Society

The Saffman-Taylor Instability Without Walls SANDRA TROIAN,
California Institute of Technology, 1200 E California Blvd, MC 128-95, Pasadena,
CA 91125, STEFFEN BERG, Shell International BV, Rijswijk, Netherlands —
The Saffman-Taylor problem represents the unstable displacement of a more vis-
cous fluid by a less viscous fluid under the action of an external pressure gradient.
Small sinusoidal deformation of the separation front gives way to repeated fingering
and tip-splitting. This repetitive process causes the transformation of an initially
featureless front to a highly ramified curve whose fractal dimension is roughly 1.7.
This instability requires that the fluids be confined by two substrates, such as in a
Hele-Shaw cell, in order to enforce the pressure gradient which drives the flow. In
this talk we describe experiments in which this instability is observed for the first
time in nanofilms freely suspended in air for which there are no confining walls. The
nanofilms consist of an aqueous surfactant solution containing hydrosoluble poly-
mer. The fractal dimension D_F ranges from $1 < D_F < 2$ and increases with the
viscosity of the bulk solution. The expanding front appears to delineate between
a surfactant-rich mobile phase and a polymer-rich less mobile phase. We describe
a phenomenological model for linearly unstable flow in which the mobility contrast
incorporates both modulation in film thickness from disjoining pressure variation as
well as the viscosity contrast from phase segregation. This extension generalizes our
understanding of this well known phenomenon.

Sandra Troian
California Institute of Technology, 1200 E California Blvd,
MC 128-95, Pasadena, CA 91125

Date submitted: 07 Aug 2010

Electronic form version 1.4