

Abstract Submitted  
for the DFD10 Meeting of  
The American Physical Society

**Sedimentation dynamics in the presence of polymer** SHMUEL M. RUBINSTEIN, Department of Physics and Harvard School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts 02138, USA, MAHESH M. BANDI, TOM KODGER, Harvard School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts 02138, USA, DAVID A. WEITZ, Department of Physics and Harvard School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts 02138, USA — We study the sedimentation of colloidal particles in polymer supplemented solution. The polymers enrich the dynamics of sedimentation by adding both particle attraction (caused by depletion interactions) and an elastic component to the flow. The sedimentation dynamics are governed by the formation, sedimentation and consequent breakup of poroelastic clusters of many particles. By making use of a custom built laser sheet microscope we are able to track Brownian one-micron particles at single particle resolution within a large ( $\sim$ cm sized) cell. This way we can resolve between bulk and boundary effects.

Shmuel M. Rubinstein  
Department of Physics and Harvard School of Engineering and  
Applied Sciences, Harvard University, Cambridge, Massachusetts 02138, USA

Date submitted: 10 Aug 2010

Electronic form version 1.4