

Abstract Submitted
for the DFD11 Meeting of
The American Physical Society

Dynamics of a laminar flow past a rotating bullet-shaped body¹

JOSÉ IGNACIO JIMÉNEZ-GONZÁLEZ, ENRIQUE SANMIGUEL-ROJAS, Universidad de Jaen, ALEJANDRO SEVILLA, Universidad Carlos III de Madrid, CARLOS MARTÍNEZ-BAZÁN, Universidad de Jaen — Numerical simulations are performed for laminar flow past a rotating bullet-shaped body of length-to-diameter ratio $L/D = 2$, ranging the non-dimensional angular rotation velocity, $\Omega = \omega R/u_\infty$, from 0 to 0.6, and covering the Reynolds number range $320 < Re < 430$. Three transition patterns stand out as Ω increases on the parametric map Ω - Re . When $\Omega \leq 0.3$, increasing rotation speeds have a stabilizing effect for the first transition, delaying the symmetry breaking bifurcation to higher values of Re , however it triggers the second bifurcation at lower Re than in the case of stationary bodies. Furthermore, the wake undergoes three different flow regimes as Re increases: axisymmetric, *frozen* (i.e. two counterrotating vortices aligned with the streamwise direction rotate around the axis with a different frequency than that of the body) and spiral flow. A different behavior has been observed in the range $0.3 \leq \Omega \leq 0.5$ where only a single transition exists. Finally, when $\Omega > 0.5$ there are also two transitions, both leading to different swirling flows.

¹Supported by the projects DPI2008-06624-C02, P07-TEP02693 and P11-TEP5702.

José Ignacio Jiménez-González
Universidad de Jaen

Date submitted: 22 Jul 2011

Electronic form version 1.4