

Abstract Submitted
for the DFD11 Meeting of
The American Physical Society

Statistics of incremental averages of passive scalar fluctuations¹

COLIN MEYER, University of California Berkeley, Department of Civil and Environmental Engineering, LAURENT MYDLARSKI, McGill University, Department of Mechanical Engineering — Whereas statistics of differences in turbulent quantities measured over a given separation have been extensively studied, statistics of the incremental averages of the same quantities (e.g., $\Sigma\theta \equiv [\theta(x+r) + \theta(x)]/2$) have only been the focus of recent research. The present work studies incremental averages of a fluctuating passive scalar (temperature) in nearly homogeneous, isotropic turbulence, generated by an active grid, with an imposed mean scalar gradient. Following the arguments of Mouri and Hori (*Phys. Fluids*, 2010) for the velocity field, we derive a scale-dependent budget for the incremental average of the scalar field fluctuations, $\Sigma\theta$. We discuss its relationship to Yaglom's four-thirds law (which pertains to differences of passive scalar fluctuations) and compare the results with the experimental data. Furthermore, the statistics of $\Sigma\theta$ are compared with those of the incremental averages of the velocity fluctuations (Σu_α).

¹Support provided by NSERC

Laurent Mydlarski
McGill University, Department of Mechanical Engineering

Date submitted: 05 Aug 2011

Electronic form version 1.4