

Abstract Submitted
for the DFD14 Meeting of
The American Physical Society

Passive appendages aid locomotion through symmetry breaking

SHERVIN BAGHERI, UGIS LACIS, Linné Flow Centre, Dept. Mechanics, KTH, Stockholm, ANDREA MAZZINO, DICCA, University of Genova, Italy, HAMID KELLAY, Université Bordeaux, France, NICOLAS BROSSE, FREDRIK LUNDELL, Linné Flow Centre, Dept. Mechanics, KTH, Stockholm, FRANCOIS INGREMEAU, Université Bordeaux, France — Plants and animals use plumes, barbs, tails, feathers, hairs, fins, and other types of appendages to aid locomotion. Despite their enormous variation, passive appendages may contribute to locomotion by exploiting the same physical mechanism. We present a new mechanism that applies to body appendages surrounded by a separated flow, which often develops behind moving bodies larger than a few millimeters. We use theory, experiments, and numerical simulations to show that bodies with protrusions turn and drift by exploiting a symmetry-breaking instability similar to the instability of an inverted pendulum. Our model explains why the straight position of an appendage in flowing fluid is unstable and how it stabilizes either to the left or right of the incoming fluid flow direction. The discovery suggests a new mechanism of locomotion that may be relevant for certain organisms; for example, how plumed seeds may drift without wind and how motile animals may passively reorient themselves.

Shervin Bagheri
Linné Flow Centre, Dept. Mechanics, KTH, Stockholm

Date submitted: 01 Aug 2014

Electronic form version 1.4