

Abstract Submitted
for the DFD16 Meeting of
The American Physical Society

Spontaneous Droplet Jump with Electro-Bouncing¹ ERIN SCHMIDT, MARK WEISLOGEL, Portland State University — We investigate the dynamics of water droplet jumps from superhydrophobic surfaces in the presence of an electric field during a step reduction in gravity level. In the brief free-fall environment of a drop tower, when a strong non-homogeneous electric field (with a measured strength between 0.39 and 2.36 kV/cm) is imposed, body forces acting on the jumped droplets are primarily supplied by polarization stress and Coulombic attraction instead of gravity. The droplet charge, measured to be on the order of $2.3 \cdot (10^{-11})$ C, originates by electro-osmosis of charged species at the (PTFE coated) hydrophobic surface interface. This electric body force leads to a droplet bouncing behavior similar to well-known phenomena in 1-g, though occurring for larger drops ~ 0.1 mL for a given range of impact Weber numbers, **We** < **20**. In 1-g, for **We** > **0.4**, impact recoil behavior on a super-hydrophobic surface is normally dominated by damping from contact line hysteresis and by air-layer interactions. However, in the strong electric field, the droplet bounce dynamics additionally include electrohydrodynamic effects on wettability and Cassie-Wenzel transition. This is qualitatively discussed in terms of coefficients of restitution and trends in contact time.

¹This work was supported primarily by NASA Cooperative Agreement NNX12A047A.

Erin Schmidt
Portland State University

Date submitted: 31 Jul 2016

Electronic form version 1.4