

Abstract Submitted
for the DNP07 Meeting of
The American Physical Society

How to Classify Three-Body Forces – and Why HARALD W. GRIESSHAMMER, Department of Physics, George Washington University — To add 3-body forces when theory and data disagree is untenable when predictions are required. For the “pion-less” Effective Field Theory at momenta below the pion-mass, I provide a recipe to systematically estimate the typical size of 3-body forces in all partial waves and orders, including external currents [1]. It is based on the superficial degree of divergence of the 3-body diagrams which contain only two-body forces and the renormalisation-group argument that low-energy observables must be insensitive to details of short-distance dynamics. Naïve dimensional analysis must be amended as the asymptotic solution to the leading-order problem depends for large off-shell momenta crucially on the partial wave and spin-combination considered. The typical strength of most 3-body forces turns out weaker than expected, demoting many to high orders. As application, the thermal cross section of $nd \rightarrow t\gamma$ bears no new 3-body force [2], besides those fixed by the triton binding energy and nd scattering length in the triton channel: $0.485(\text{LO}) + 0.011(\text{NLO}) + 0.007(\text{NNLO})$; $\text{mb} = [0.503 \pm 0.003] \text{ mb}$, converges and compares well with data, $[0.509 \pm 0.015] \text{ mb}$. Potential models list $[0.49 \dots 0.66] \text{ mb}$, depending on the 2-nucleon potential and inclusion of the $\Delta(1232)$. [1] H.W. Griesshammer: Nucl. Phys. **A760** (2005) 110 [2] H. Sadeghi, S. Bayegan and H.W. Griesshammer: Phys. Lett. **B643** (2006), 263.

Harald W. Griesshammer
Department of Physics, George Washington University

Date submitted: 26 Jun 2007

Electronic form version 1.4