

Abstract Submitted
for the DNP15 Meeting of
The American Physical Society

Nuclear Spin Dependent Parity Violation in Diatomic Molecules

SIDNEY CAHN, EMINE ALTUNTAS, Yale University, JEFFREY AMMON, MIT Lincoln Labs, DAVID DEMILLE, Yale University — Nuclear spin-dependent parity violation (NSD-PV) effects arise from the exchange of the Z^0 boson between electrons and the nucleus and from the interaction of electrons with the nuclear anapole moment, a parity-odd magnetic moment. The anapole moment grows as $A^{2/3}$ of the nucleus, while the Z^0 coupling is independent of A . We study NSD-PV effects using diatomic molecules, where signals are dramatically amplified by bringing rotational levels of opposite parity close to degeneracy in a strong magnetic field. Using a Stark-interference technique, we measure the NSD-PV interaction matrix element. We present results that demonstrate statistical sensitivity to NSD-PV effects surpassing that of any previous atomic parity violation measurement, using the test system $^{138}\text{Ba}^{19}\text{F}$. We also discuss improvements on investigations of systematics due to non-reversing stray E -fields, E_{nr} together with B -field inhomogeneities, and short-term prospects for measuring the nuclear anapole moment of ^{137}Ba . In the long term, our technique is sufficiently general and sensitive to enable measurements across a broad range of nuclei.

Sidney Cahn
Yale University

Date submitted: 09 Jul 2015

Electronic form version 1.4