

Abstract Submitted
for the DNP16 Meeting of
The American Physical Society

A Study on Low Spin States in ^{154}Gd Using (p,p') Reaction¹

HARRIS BIDAMAN, B. HADINIA, P. E. GARRETT, University of Guelph, UNIVERSITY OF LIVERPOOL COLLABORATION, DARESBUURY LABORATORY COLLABORATION, UNIVERSITY OF JYVÄSKYLÄ COLLABORATION, UNIVERSITY OF GUELPH COLLABORATION — Located at the stability line, the low lying spin states of the ^{154}Gd nucleus were investigated at the University of Jyväskylä accelerator laboratory in Finland using the $^{154}\text{Gd}(p,p'\gamma)$ reaction. A proton beam of 12 MeV was used to excite the ^{154}Gd target, with the gamma-rays from the reaction detected with the JUROGAM II array, while the LISA charged-particle spectrometer was used for detection of the inelastically scattered protons. This experiment marked one of the first uses of the LISA spectrometer at Jyväskylä, which enabled the efficient tagging of the proton-emitting reactions, thus helping to distinguish between the (p,p') and the much more copious (p,xn) channels. By analysing the peaks obtained from the gamma-gamma, and gamma-gamma-proton coincidence matrices, a decay scheme has been built using the RadWare software Esc18r. Experimental methods, new transitions, and future steps will be discussed.

¹Canada Foundation for Innovation, the Natural Sciences and Engineering Research Council of Canada, the National Research Council of Canada and the Canadian Research Chairs program.

Harris Bidaman
University of Guelph

Date submitted: 10 Aug 2016

Electronic form version 1.4