

Abstract Submitted
for the DNP19 Meeting of
The American Physical Society

Study of Kaon Identification for Hyperon Photoproduction in JLab's Hall B CLAS12¹ ISABELLA ILLARI, WILLIAM PHELPS, George Washington University, CLAS COLLABORATION — The CLAS12 physics program is involved in the study of baryon spectroscopy in quasi-real photoproduction of a large variety of final states, and includes the photoproduction of singly, doubly and triply strange hyperons. The photoproduction of the very strange hyperon, the Ω^- , is of particular interest. Its photoproduction cross section is unknown. Furthermore, the dynamics of the Ω^- photoproduction is unclear, as there are no strange quarks in the initial state while there are three in the final state. The available theoretical predictions for the Ω^- hyperon photo- and electroproduction cross section vary from 1 to 300 pb. As of today, there is only an upper limit of the cross section ($\sigma_{tot} < 17$ nb at 20 GeV) reported by SLAC. To identify the final state for $\gamma p \rightarrow \Omega^- K^+ K^+ K^0$ the minimal requirement is to detect three kaons: two K^+ and one K_s^0 . The latter is identified by its decay to $\pi^+ \pi^-$. We will discuss our study of kaon identification in CLAS12 based on Monte Carlo simulations and real data collected by Run Group A, which used a ~ 11 GeV beam incident on a liquid hydrogen target.

¹This work was performed with partial support from US DOE DE-SC001658, The George Washington University, the CLAS Collaboration, and Thomas Jefferson National Accelerator Facility.

Isabella Illari
George Washington University

Date submitted: 30 Jun 2019

Electronic form version 1.4