

Abstract Submitted
for the DNP20 Meeting of
The American Physical Society

Dielectron production in Au+Au collisions at $\sqrt{s_{NN}} = 54$ GeV at STAR ZHEN WANG, Shandong Univ, STAR COLLABORATION — Dielectrons are excellent probes of the Quark-Gluon Plasma (QGP) created in high-energy heavy-ion collisions. Because they can be produced at all stages of the collision system evolution and do not interact with the medium strongly, which indicates that dielectrons carry the information from the initial stage to the final stage. In the low mass region (LMR, $M_{ee} < M_\phi$), the mass spectra of vector mesons will be modified by the hot and dense medium which is related to the chiral symmetry restoration in the medium. In the intermediate mass region (IMR, $M_\phi < M_{ee} < M_{J/\psi}$), QGP thermal radiation can be used as a QGP thermometer. However, it is complicated to measure the QGP thermal radiation because of the heavy flavor semi-leptonic decay contributions. In this presentation, I will present the dielectron production in Au+Au collisions at $\sqrt{s_{NN}} = 54$ GeV at STAR. With a 10 times larger data sample than that at 62 GeV from the first phase of the STAR Beam Energy Scan (BES-I) program, in-medium ρ modification can be studied with better precision and compared to different theoretical predictions. Furthermore, the physics implications of these measurements will be discussed.

Zhen Wang
Shandong Univ

Date submitted: 24 Jun 2020

Electronic form version 1.4