

Abstract Submitted
for the DNP20 Meeting of
The American Physical Society

Constraining the Sea Quark Distributions Through W and Z Cross Sections and Cross-Section Ratios Measured at STAR¹ MATTHEW POSIK, Temple Univ, STAR COLLABORATION — Although parton distribution functions (PDFs) have become more precise, there are still kinematic regions where more data are needed to help constrain global PDF extractions, such as the ratio of the sea quark distributions \bar{d}/\bar{u} near the valence region. Furthermore, different measurements appear to suggest different high- x behaviors of this ratio. The W cross-section ratio (W^+/W^-) is sensitive to the unpolarized quark distributions at large Q^2 . Such a measurement can be used to help constrain the \bar{d}/\bar{u} ratio. The STAR experiment at RHIC is well equipped to measure the leptonic decays of W and Z bosons, in the pseudorapidity range ($-1.0 < \eta < 1.5$), produced in proton-proton collisions at $\sqrt{s} = 500/510$ GeV. These cross sections and their ratios are sensitive to quark and antiquark distributions in the x -range $0.1 < x < 0.3$. This talk will present preliminary results from the 2011-2013 RHIC runs, which total about 350 pb^{-1} of integrated luminosity. Measurements of the differential W and Z cross sections and the W^+/W^- cross-section ratio as a function of the decay lepton's pseudorapidity, the W/Z cross-section ratio and total W and Z cross sections will be shown.

¹DOE NP contract: DE-SC0013405

Matthew Posik
Temple Univ

Date submitted: 26 Jun 2020

Electronic form version 1.4